

Published in final edited form as:

Cell Microbiol. 2014 August ; 16(8): 1146–1155. doi:10.1111/cmi.12312.

Myeloperoxidase in human neutrophil host defense

William M. Nauseef¹

¹Inflammation Program, Department of Medicine, Roy J. and Lucille A. Carver College of Medicine, University of Iowa, and Veterans Administration Medical Center, Iowa City, IA 52242

Background

While in search of a tissue source of catalase that would lack the biliverdin that contaminated liver preparations, Kjell Agner discovered in tuberculous empyema a green enzyme that was structurally different from other proteins known at that time (Agner, 1941). Agner described many of the structural and enzymatic properties of the green enzyme, verdoperoxidase, in his doctoral thesis for medical school (Agner, 1941). Two years later, in 1943, Theorell and Åkeson purified a brown-green peroxidase from cow's milk and demonstrated that it had spectral properties distinct from those of Agner's leukocyte-derived verdoperoxidase (Theorell et al., 1943). Recognizing that despite both enzymes being green, they originated from different sources, Theorell and Åkeson proposed the names myeloperoxidase for Agner's enzyme and lacto-peroxidase for the milk enzyme.

Whereas Agner's subsequent studies detailed how myeloperoxidase (MPO) inactivated bacterial toxins (Agner, 1941; Agner, 1950; Agner, 1955), Schultz initiated seminal studies to define the biological relevance of MPO in a broader context. Noting that MPO represented >5% of the dry weight of human neutrophils (Schultz et al., 1962), Schultz purified the protein from neutrophils isolated from human blood and defined many of the structural and enzymatic features of MPO. Pioneering studies by Zgliczynski, Schultz and others elucidated much of the chemistry supported by MPO and speculated how it might participate in human host defense (Schultz et al., 1962; Zgliczynski et al., 1968; Zgliczynski et al., 1971; Harrison, et al., 1976; Zgliczynski et al., 1980). Seminal studies by Klebanoff and colleagues established the critical links between the presence of MPO in neutrophils, halogenation of ingested bacteria in phagosomes, and the microbicidal action of the MPO-H₂O₂-Cl system (Klebanoff, 1967; Klebanoff, 1968; Klebanoff, 1970a; Klebanoff, 1970b; Klebanoff et al., 1972; Klebanoff, 2005).



Based on a comprehensive analysis of the evolutionary relationships among sequences of peroxidase proteins in the plant and animal kingdoms, the Obinger laboratory has recently categorized MPO and other proteins previously called animal peroxidases as members of Chordata peroxidase subfamily (Zamocky *et al.*, 2008). All members expressed in mammals, including MPO, EPO, LPO, and TPO, contain conserved motifs on both proximal and distal sides of the essential heme prosthetic group, a calcium-binding site, and at least

Correspondence: William M. Nauseef, Inflammation Program, 2501 Crosspark Road, Coralville, IA, USA 52241. Phone (319) 335-4278, Fax (319) 335-4194; william-nauseef@uiowa.edu.

Disclosures: The author has nothing to disclose.

two covalent bonds linking the heme group to the protein backbone. In addition to the conserved ester linkages in the heme pocket, MPO contains a third, a sulfonium linkage between the 2-vinyl group and methionine 409. The presence of the sulfonium linkage empowers MPO with unique spectral qualities as well as greater oxidizing potential, rendering MPO the only member of the protein superfamily able to oxidize Cl^- to Cl^+ and thereby generate HOCl at physiologic pH (Furtmuller *et al.*, 2003; Arnhold *et al.*, 2006; Zederbauer *et al.*, 2007).

Collectively, all peroxidases support the same general chemistry; they catalyze one- and two-electron oxidations of susceptible substrates, both organic (AH_2) and inorganic, such as halides (X^-), to promote structural changes with physiological consequences:



Peroxidase-mediated post-translational modifications underlie a diverse set of biological phenomena, from thyroid hormone synthesis (Ruf *et al.*, 2006.) to pathophysiologic lipoprotein modification (Undurti *et al.*, 2009), to crosslinking of fertilized sea urchin eggs to block entry of sperm arriving too late (Foerder *et al.* 1977). In all situations, the principles remain the same, only the substrate and context differ. This review focuses attention on MPO-mediated events in neutrophil phagosomes, the highly specialized space where ingested microbes are confined, and in the context of their contribution to host defense against infection. Extensive and elegant studies have examined MPO-dependent biochemistry in other inflammatory diseases, most notably cardiovascular disease, and the interested reader will be richly rewarded by reading any of several recent and excellent reviews (Malle *et al.*, 2003; Nicholls *et al.*, 2009; Shao *et al.*, 2012; Nussbaum *et al.*, 2013).

Players

The essential elements of heme peroxidase-based biochemistry include the peroxidase, a source for H_2O_2 , and substrates that can undergo one or two electron oxidation (Dunford, 1999). In the context of events within the phagosome of human neutrophils, the identities of the peroxidase and the H_2O_2 source, MPO and the phagocyte NADPH oxidase, respectively, are well established (Figure). In addition, the voltage-gated proton channel, Hv1, provides the charge compensation required to sustain NADPH oxidase activity (DeCoursey, 2013). Less well characterized is the source of chloride to support HOCl production. The MPO- H_2O_2 -Cl system requires a continuous supply of Cl, as the small amounts internalized incidentally during phagocytosis are rapidly consumed as degranulation and NADPH oxidase activation occur (Winterbourn *et al.*, 2006). The unusually high chloride concentration in neutrophil cytoplasm provides a reservoir to meet intraphagosomal needs, and several different transporters serve to deliver the needed chloride. Cystic fibrosis transmembrane conductance regulator (CFTR) transports the bulk of the chloride from cytoplasm into the phagosomal lumen (Aiken *et al.*, 2012; Zhou *et al.*, 2013). CFTR, a constituent of secretory vesicle, but not plasma, membrane in resting neutrophils, reaches

the nascent phagosomes as granules fuse during phagocytosis, thereby providing the means to transfer chloride from the cytoplasmic reserves (Painter *et al.*, 2006; Zhou *et al.*, 2013). The steady state concentrations of chloride in phagosomes of patients with cystic fibrosis [25.5 ± 3.3 mM, $n = 5$] or normal neutrophils treated with the CFTR inhibitor GlyH-101 [27.7 ± 4.4 mM, $n = 5$] are significantly lower than that in normal neutrophil phagosomes [67.7 ± 7.3 mM, $n = 5$] (Painter *et al.*, 2010). Although the measured values may be higher than the *bona fide* steady state concentration due to technical reasons¹, optimal MPO activity in phagosomes requires chloride transport via CFTR; phagosomes containing genetically or pharmacologically dysfunctional CFTR exhibit reduced MPO-dependent modification of targets (*vide infra*) and killing of ingested *Pseudomonas aeruginosa* (Painter *et al.*, 2008). In addition to CFTR, the anion transporter ClC3 (Moreland *et al.*, 2006) and the K^+/Cl^- co-transporter KCC3 (Sun *et al.*, 2012.) also contribute to chloride transport into neutrophil phagosomes.

Products of the MPO-H₂O₂-Cl system

Compound I, the Fe(IV) cation radical produced by the reaction between H₂O₂ and native MPO (Marquez *et al.*, 1994; Furtmüller *et al.*, 1998), can support one-and two-electron oxidations of an array of substrates, including halides, the pseudohalide thiocyanate, amino acids, proteins, lipids, and nucleic acids [reviewed comprehensively in (Pattison *et al.*, 2006; Summers *et al.*, 2008)]. Even after restricting consideration to the chemistry plausible in the phagosome, the number of potential reactions exceeds facile comprehension. Rather than catalogue the multiple and diverse possibilities, I will summarize key reactions likely to occur in human neutrophils and contribute, directly or indirectly, to killing ingested organisms. Many of the details of reactions and their kinetics will be omitted here, instead directing the interested reader to recently published excellent reviews (Hurst, 2012; Klebanoff *et al.*, 2013; Winterbourn *et al.*, 2013).

For the sake of simplifying the presentation, consider HOCl as the most proximal effector product of the MPO-H₂O₂-Cl system. The product of the two-electron oxidation of Cl^- , HOCl exists in equilibrium with OCI^- at the near neutral pH in neutrophil phagosomes, as the pK_a is 7.53 at 30°C. Under conditions that include substrates that would compete with Cl^- for oxidation by Compound I, the number of potential reaction substrates multiplies. The reactions of HOCl with amino acids, peptides, and proteins have been extensively studied [review (Hawkins *et al.* 2003)] and generate products, including chloramines and eventually aldehydes (Zgliczynski *et al.*, 1971) that can exert antimicrobial activity as well (Thomas *et al.*, 1986), and can generate tyrosyl radicals that prompt protein dimerization and aggregation (Vissers *et al.*, 1991). HOCl generated by MPO produces mono- and dichloramines of primary amines, including those on neutrophil granule proteins present within phagosomes (Thomas *et al.*, 1982). Although these represent quantitatively minor reactions, they can be long-lived and, depending on the concentration and structural features, can contribute to antimicrobial action. Some products, such as the monochloramine NH₂Cl

¹Note that these measurements likely overestimate the true chloride concentration, because the intrinsic susceptibility of the fluorescent probe used to detect chloride requires inclusion of sodium azide in experimental conditions. Consequently, chloride consumption by MPO is inhibited under the experimental conditions.

generated by the reaction of HOCl and NH_4^+ , can penetrate the hydrophobic surfaces of particular microbes and thereby gain access to vulnerable intracellular targets (Thomas, 1979). In the case of *E.coli*, MPO-dependent generation of long-lived chloramines and chloramide derivatives correlates with oxidation of critical bacterial sulfhydryl groups and loss of microbial viability (Thomas, 1979a; Thomas, 1979b). The magnitude of the contribution of chloramines and other HOCl-derived agents to antimicrobial action varies as a function of the size, charge, and polarity of the toxin produced, the chemical composition of the surface of the ingested microbe, and the specific vulnerable targets in the organism.

The abundant supply of MPO ideally equips neutrophils for service in innate immune defense against microbes, converting the H_2O_2 produced by the NADPH oxidase into the potent microbicide HOCl. H_2O_2 possesses greater redox potential than does HOCl (1.776V vs 1.482V) but exerts much less antimicrobial action (Klebanoff, 1978; Hurst *et al.*, 1989; Elzanowska *et al.*, 1995). For example, *E.coli* washed with EDTA to remove adventitious metal cations that might support Fenton chemistry can tolerate 50 mM H_2O_2 for more than 40 minutes without a loss in viability (Elzanowska *et al.*, 1995). In part, this tolerance reflects the detoxifying action of peroxide-consuming enzymes present in aerobic organisms, as without such enzymes, even endogenously produced H_2O_2 can exert substantial DNA damage (Park *et al.*, 2005). H_2O_2 oxidizes cysteine and methionine residues most readily but at relatively slow rates compared to the activity of HOCl. For example, the rate constant for thiol oxidation by H_2O_2 is $2.9 \text{ M}^{-1}\text{s}^{-1}$ (Imlay, 2003), whereas that for HOCl is $\sim$ a million-fold faster, $3.0 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$ (Pattison *et al.*, 2001). The rapid reaction kinetics of HOCl enhances its capacity to attack microbial targets that may be accessible only briefly. Supporting that prediction, HOCl, but not H_2O_2 , prompts protein aggregation *in vitro* (Winter *et al.*, 2008), presumably by irreversibly modifying vulnerable targets that are exposed only transiently during the continuous folding and unfolding of protein conformations.

Targets of the MPO- H_2O_2 -Cl system

The specific target and the HOCl-mediated changes that culminate in microbial death have not been defined and vary among organisms (Hurst, 2012). All things being equal with respect to inherent susceptibility of microbial components to modification, spatial arrangements dictate that surface structures would be the first to encounter MPO-derived toxins. However, in many cases, HOCl attacks surface structures but with little consequence on bacterial viability. Instead, loss of function supported by proteins in bacterial inner membranes correlates with microbial death [reviewed in (Hurst, 2012)], as does HOCl-dependent oxidation of methionine residues in cytosolic and inner membrane proteins of *E.coli* (Rosen *et al.*, 2009). Whatever the underlying chemistry, HOCl kills within milliseconds (Albrich *et al.*, 1982) and more efficiently than does H_2O_2 in a variety of experimental settings. Based on the number of molecules of oxidant/organism needed to kill *E.coli*, HOCl is $\sim$ 1000-fold more potent than is H_2O_2 ($0.4\text{--}5.0 \times 10^8$ vs $>3 \times 10^{11}$) (Lymer *et al.*, 1995). Similarly, the relative dose-dependent killing of *K.pneumoniae* demonstrates 50% reduction in viability at 60 min for 1 μM HOCl vs 1 mM H_2O_2 (Hirche *et al.*, 2005). Like many experimental systems, both of these examples utilized buffer systems without protein and were thus free of competing substrates for the oxidants. As discussed later, teasing out

“lethal hits” from the array of competing reactions with inadvertent targets that constitute complex biological systems, as in neutrophil phagosomes, represents a formidable challenge to constructing *in vitro* conditions that accurately mirror event in neutrophils and be used to assess how neutrophils kill.

Diversity among the microbes with respect to surface composition (*e.g.* presence or absence of capsule, composition of capsule if present, lipopolysaccharides in enteric Gram negative rods vs lipoteichoic acid in Gram positive bacteria) undermine the notion of the existence of a single microbial target. Furthermore, microbes sense attack by antimicrobial agents and respond to threats to their viability by evading toxins, enduring the modifications to structure or function, repairing host-induced damage, or all of the aforementioned actions. In some cases, the bacterial responses match predictions based on current understanding of events in phagosomes. The different transcriptional responses of *E.coli* ingested by neutrophils from normal individuals vs those with chronic granulomatous disease (CGD), an inherited deficiency in NADPH oxidase activity, illustrate the remarkable specificity and speed with which an organism can respond to its environment. For example, *E.coli* fed to normal neutrophils modulate expression of > 70 genes, including upregulation of genes such as *OxyRS* that respond to H₂O₂ (Staudinger *et al.*, 2002). In contrast, the same *E.coli* fed to CGD neutrophils failed to upregulate H₂O₂-responsive genes, as anticipated given the absence of oxidants in CGD phagosomes. In other settings, bacterial responses affirm the complexity of biochemistry in normal neutrophils. MPO-mediated HOCl oxidizes methionine residues, and methionine sulfoxide (MeO) production correlates with loss of viability of *E.coli* (Rosen *et al.*, 2009). Deletion of methionine sulfoxide reductase (*msr*) from *E.coli* eliminates the capacity to regenerate methionine from MeO and increases susceptibility to HOCl. Likewise, *msr*-deletion renders *S.aureus* more susceptible to oxidant and neutrophil killing (Pang *et al.*, 2014.). These data jibe with expectations, given the recognized importance of oxidants in killing and the role of *msr* in the repair of oxidant damage. Whereas *S.aureus* ingested by normal neutrophils increase expression of *msr*, upregulation is far greater in neutrophils without NADPH oxidase activity. Furthermore, purified neutrophil granules alone, without added oxidants, promote *msr* upregulation in *S.aureus*. Collectively, these data demonstrate that ingested bacteria react promptly and selectively to neutrophil-derived toxins but often in ways not readily explained by current models of antimicrobial activity. However, the unexpected observation that *S.aureus* upregulates *msr* under conditions free of exogenous oxidants illustrates the complex interactions among phagosomal contents.

In situ

A significant fraction of the oxygen consumed by activated neutrophils can be recovered as HOCl, and a sensitive, specific, and readily performed assay can be used to quantitate extracellular HOCl production by stimulated neutrophils (Dypbukt *et al.*, 2005). HOCl represents from 28 to 72% of the oxygen consumed by phorbol myristate acetate (PMA)-stimulated neutrophils (Weiss *et al.*, 1982; Foote *et al.*, 1983; Thomas *et al.*, 1983; Chapman *et al.*, 2002). The susceptibility of the fluorophore green fluorescent protein (GFP) to bleaching by chlorination of Y66 (Espey *et al.* 2002) provides a specific and selective tool to monitor HOCl generation with neutrophil phagosomes. Both *E.coli* (Palazzolo *et al.* 2005)

and *S.aureus* (Schwartz *et al.*, 2009) stably expressing GFP have served as probes for neutrophil HOCl production. In addition to HOCl being generated within phagosomes, the loss of viability directly correlates with MPO-dependent modification of ingested bacteria.

Compartmentalization of ingested microorganisms within phagosomes not only limits their access to vital nutrients but also creates conditions that enhance toxicity of antimicrobial agents present. From the perspective of chemistry, the close approximation of phagosomal membrane to the microbial surface creates a relatively small reaction zone for oxidants to attack soluble substrates in phagosomal fluid. Consequently, the tight quarters favor reactions between oxidants and microbe even in an environment rich in competing substrates (Lyman *et al.* 1995). Additionally, the very small volume of the phagosome, estimated to be 1.2 fl (Winterbourn *et al.*, 2006), allows high concentrations of oxidants to be achieved, and the ability of MPO to adhere to some bacteria enhances the effective concentration of toxic agents at the site of attack.

Adopting a reductionist approach to examine *in vitro* the antimicrobial activity of MPO, most investigators utilize the MPO-H₂O₂-Cl system (Klebanoff, 1968) free of other neutrophil granule proteins. However, during phagocytosis, fusion of neutrophil granules delivers agents, including proteolytic enzymes and proteins with direct antimicrobial activity [reviewed in (Rorvig *et al.*, 2013)], that contribute to the antimicrobial activity and can serve as substrate for HOCl. Accessibility and inherent susceptibility determine sites for attack by the MPO-H₂O₂-Cl system, rendering both exogenous substrates originating from ingested microbes and endogenous neutrophil-derived molecules released into phagosomes during degranulation equally eligible targets.

The complex mix of synergistic as well as antagonistic interactions dictates the fate and overall survival of bacteria in neutrophils (Nauseef, 2007; Hurst, 2012; Winterbourn *et al.*, 2013). Neutrophil elastase (NE) enhances the killing of *S.aureus* as well as *E.coli* by MPO-H₂O₂-Cl by a mechanism that is independent of protease activity, as heating NE does not compromise its synergistic influence (Odeberg *et al.*, 1976). NE augments cathepsin G-mediated killing as well, but cathepsin G does not synergize with the MPO-H₂O₂-Cl system. Adding to the complexity of so many interactions among effector molecules, MPO-mediated modifications can inactivate many of the granule proteins in neutrophils (Voetman *et al.*, 1981). Although only some have been examined, many granule proteins undergo irreversible inactivation in the presence of HOCl or MPO-H₂O₂-Cl, including acid phosphatase, N-acetyl- β -glucosaminidase, β -glucuronidase, α -fucosidase, α -mannosidase, lysozyme, vitamin B₁₂-binding protein, MMP-7, -8, and -9, cathepsin G, and NE (Kobayashi *et al.*, 1982; Clark *et al.*, 1985; Fu *et al.*, 2001; Fu *et al.*, 2003; Hawkins *et al.*, 2005; Hirche *et al.*, 2005). Whereas HOCl or the MPO-H₂O₂-Cl system inactivates NE, cathepsin G, and MMP-7, H₂O₂ alone does not (Fu *et al.*, 2003; Hirche *et al.*, 2005; Shao *et al.*, 2005). Furthermore, HOCl at low concentrations inactivates NE without prompting changes in apparent size after SDS-PAGE, whereas at high concentrations of HOCl NE undergoes complete degradation (Hirche *et al.*, 2005), suggesting that extensive oxidation of NE by HOCl may induce its autodegradation.

MPO can mediate the same inactivation of lysosomal enzymes in more complex biological settings, as in supernatants from stimulated neutrophils. Supernatants from stimulated neutrophils from patients with chronic granulomatous disease or MPO deficiency, cells lacking essential elements for MPO-mediated chemistry, contain more lysosomal enzyme activity than do secretions from normal neutrophils, and the activity of lysosomal enzymes released by normal neutrophils are increased under anaerobic conditions, in the presence of inhibitors of MPO or the NADPH oxidase, or by adding agents that scavenge oxidants (Kobayashi *et al.*, 1982; Clark *et al.*, 1985; Weiss *et al.*, 1985; Fu *et al.*, 2003; Hirche *et al.*, 2005). Not directly relevant to host-pathogen interactions but pertinent to inflammation in general, MPO can oxidatively inactivate plasma protease inhibitors and thereby modulate extracellular tissue damage at inflammatory sites (Clark *et al.*, 1979; Clark *et al.*, 1980; Clark *et al.*, 1981; Clark, 1983; Hawkins *et al.*, 2005). As a result of egalitarian attack of the MPO-H₂O₂-Cl system on substrates whether from man or microbe, the granule proteins released into phagosomes can be chemically modified and functionally modulated, enhanced as well as reduced, depending on the particular agent and the specific modification. Recognition of such interactions and their consequences figures into the final analysis of overall events in the neutrophil phagosome.

Role for MPO in the antimicrobial activity of intact neutrophils

Neutrophils possess MPO and can generate HOCl, which can exert rapid microbicidal action on a wide variety of microorganisms, but do they rely on MPO to kill ingested microbes trapped in phagosomes?

Studied *in vitro* and compared to normal neutrophils, those deficient in MPO have defective killing of bacteria as well as fungi, with variation among organisms in the degree of the defect. For example, killing of *S.aureus* after 30 minutes by MPO-deficient neutrophils is ~30% of that by normal neutrophils but ~80% of normal when *E.coli* are fed to neutrophils (Klebanoff *et al.*, 1972; Klebanoff *et al.*, 2013). Furthermore, MPO-deficient neutrophils kill susceptible bacteria more slowly than do normal neutrophils, but in some cases achieve the same success in the end (Lehrer *et al.*, 1969; Hampton *et al.*, 1996; Decleva *et al.*, 2006). In contrast to the slow but eventually effective killing of selected bacteria, MPO-deficient neutrophils lack the capacity to kill *Candida albicans* (Decleva *et al.*, 2006), a defect that parallels the clinical picture of some individuals with MPO deficiency (*vide infra*). The variation in antimicrobial activity as a function of the specific organism is not unique to MPO deficiency but seen as well in chronic granulomatous disease (Holland, 2013). In fact, normal, CGD, and MPO-deficient neutrophils kill *E.coli* equally well (Rosen *et al.*, 1997).

Experimental infection models using MPO knock-out mice have demonstrated increased susceptibility of deficient mice to challenges with a variety of organisms, including *Klebsiella* and *Candida* (Aratani *et al.*, 1999; Aratani *et al.*, 2000; Hirche *et al.*, 2005; Aratani *et al.*, 2006), and to the cecal ligation and puncture model (Gaut *et al.*, 2001). Data from some of the animal studies highlight a relationship between organism load and the impact of MPO on outcome of infection. When the inoculum of infection with fungi is high, eradication of infection and survival require MPO. In contrast, oxidants generated by the NADPH oxidase suffice in responding effectively to challenges with low inocula (Aratani *et*

et al., 2002a; Aratani *et al.*, 2002b). Although the validity of extrapolations from MPO knock-out mice to humans can be undermined by inherent differences between phagocytes from the two species with respect to NADPH oxidase activity, MPO content, and the repertoire of MPO-independent antimicrobial agents such as defensins (Rausch *et al.*, 1975; Eisenhauer *et al.*, 1992; Mestas *et al.*, 2004; Rittirsch *et al.*, 2007), the contributions of MPO to the inflammatory process *per se*, independent of direct antimicrobial action, also obfuscate interpretation of the results of experimental models. MPO knock-out animals express higher levels of induced nitric oxide synthase, generate more nitric oxide, and exhibit less lung injury in response to sepsis or endotoxin (Brovkovych *et al.*, 2008; Haegens *et al.*, 2009; Takeuchi *et al.*, 2012), demonstrating that overall survival from infection reflects important factors in addition to the eradication of viable organisms.

As reviewed recently (Klebanoff *et al.*, 2013), widespread application of flow cytometry in clinical laboratories upended appreciation of the prevalence and clinical phenotype of inherited MPO deficiency in humans. Only two patients with MPO deficiency had been reported (Grignaschi *et al.*, 1963; Undritz, 1966) prior 1969 when Lehrer and Cline described a young diabetic male with MPO deficiency and disseminated candidiasis (Lehrer *et al.*, 1969) and reports of MPO deficiency were infrequent until hematology laboratories adopted peroxidase staining and flow cytometry to identify and enumerate cells in peripheral blood. Subsequently, studies demonstrate that MPO deficiency occurs relatively commonly, affecting 1 in 2,000 to 4,000 healthy individuals in North America and Europe (Kitahara *et al.*, 1981; Kutter *et al.*, 1994; Kutter, 1998) and 1 in 57,000 in Japan (Nunoi *et al.*, 2003). With the exception of systemic candidiasis in the presence of diabetes mellitus, few MPO-deficient patients with increased or unusual infections have been reported. Although the paucity of infectious complications in MPO-deficient subjects could be offered as evidence against the importance of MPO in host response to infection, that would be a superficial interpretation of the data, as MPO-deficient neutrophils are not equivalent to neutrophils that lack only MPO (Klebanoff, 1970). Klebanoff demonstrated that MPO-deficient neutrophils kill test microbes more efficiently than do normal neutrophils treated with azide (to inhibit MPO) and that killing by MPO-deficient neutrophils is not inhibited by azide treatment, together suggesting that myeloid cells in MPO-deficient subjects may adapt to the deficiency by enlisting antimicrobial systems that compensate. However, the absence of MPO has consequences for antimicrobial action that reflect the loss of HOCl-dependent modification of neutrophil granule proteins.

The many synergies among antimicrobial toxins acting in phagosomes sabotage reaching a straightforward answer to questions about the role of MPO in human host defense. The absence of MPO eliminates not only the capacity to deliver lethal damage to target microbes but also to modify the structure and activity of other antimicrobial elements present in phagosomes. Without MPO, granule proteins such as NE, cathepsin G, and MMP-7, agents otherwise inactivated by the MPO-H₂O₂-chloride system, will be functional and exert antimicrobial action not present in the phagosomes of normal neutrophils. Furthermore, the ambient H₂O₂ concentration in phagosomes lacking MPO will be higher than that in normal neutrophils for two reasons. MPO terminates activity of the NADPH oxidase (Jandl *et al.*, 1978) and MPO-deficient neutrophils have prolonged oxidase activity and generate more

H₂O₂ than do normal neutrophils (Klebanoff *et al.*, 1971; Klebanoff *et al.*, 1972; Rosen *et al.*, 1976; Nauseef *et al.*, 1983). In addition, the absence of MPO, which normally rapidly consumes H₂O₂ to generate HOCl, will further increase the concentration of H₂O₂ (Winterbourn *et al.*, 2006). The altered oxidant composition and tone in MPO-deficient phagosomes modifies not only the available effector molecules but also substrates on and in microbes that constitute vulnerable targets. Microbial sites may be more or less susceptible to attack after modification by H₂O₂ vs HOCl. Taken together, these factors suggest that contrasting the prevalence of infection *in vivo* or antimicrobial action *in vitro* of normal and MPO-deficient neutrophils overlooks the downstream effects of MPO-H₂O₂-Cl on the collective antimicrobial activity of other oxidants and granule proteins in phagosomes and offers only limited insight into the contribution of MPO to normal neutrophil function.

Take Home

As noted nearly four decades ago, initial neutrophil-dependent killing of microbes depends on a system that can be inhibited by azide (Koch, 1974). The most rapid and complete antimicrobial action by human neutrophils against many organisms relies on H₂O₂ from the phagocyte oxidase and MPO from azurophilic granules. MPO supports production of lethal amounts of HOCl in human neutrophils, bacteria recovered from phagosomes have MPO-specific posttranslational modifications, MPO-mediated changes in bacteria correlate with loss of their viability, and MPO-deficient neutrophils kill many species of microorganisms more slowly than do normal cells. Collectively, these data suggest that efficient antimicrobial activity by neutrophils depends on the MPO-H₂O₂-Cl system.

The strikingly different clinical phenotypes of humans lacking the NADPH oxidase or MPO highlights the complexity of interactions within phagosomes that culminate in the death of ingested microbes. The dominant oxidant present in normal (HOCl), MPO-deficient (H₂O₂), and CGD (none) neutrophils react differently with both microbial targets as well as vulnerable sites on granule proteins, thereby altering the overall antimicrobial “tone” in phagosomes in the distinctly different settings.

Acknowledgments

Supported by grants AI070958 and AI044642 from the National Institutes of Health (WMN). The Nauseef lab is also supported by a Merit Review award and use of facilities at the Iowa City Department of Veterans Affairs (VA) Medical Center, Iowa City, IA, USA 52246.

References

- Agner K. Verdoperoxidase: a ferment isolated from leukocytes. *Acta Physiol Scand.* 1941; 2:1–62.
- Agner K. Detoxicating effect of verdoperoxidase on toxins. *Nature.* 1947; 159:271–272. [PubMed: 20287530]
- Agner K. Studies on peroxidative detoxification of purified diphtheria toxin. *J Exp Med.* 1950; 92:337–347. [PubMed: 14778915]
- Agner K. Peroxidative detoxification of diphtheria toxin studied by using I 131. *Rec Trav Chim.* 1955; 74:373–376.
- Aiken ML, Painter RG, Zhou Y, Wang G. Chloride transport in functionally active phagosomes isolated from Human neutrophils. *Free Radic Biol Med.* 2012; 53:2308–2317. [PubMed: 23089227]

- Albrich JM, Hurst JK. Oxidative inactivation of *Escherichia coli* by hypochlorous acid. Rates and differentiation of respiratory from other reaction sites. *FEBS Lett.* 1982; 144:157–161. [PubMed: 6286355]
- Aratani Y, Koyama H, Nyui SI, Suzuki K, Kura F, Maeda N. Severe impairment in early host defense against *Candida albicans* in mice deficient in myeloperoxidase. *Infect Immun.* 1999; 67:1828–1836. [PubMed: 10085024]
- Aratani Y, Kura F, Watanabe H, Akagawa H, Takano Y, Ishida-Okawara A, et al. Contribution of the myeloperoxidase-dependent oxidative system to host defense against *Cryptococcus neoformans*. *J Med Microbiol.* 2006; 55:1291–1299. [PubMed: 16914663]
- Aratani Y, Kura F, Watanabe H, Akagawa H, Takano Y, Suzuki K, et al. Critical role of myeloperoxidase and nicotinamide adenine dinucleotide phosphate-oxidase in high-burden systemic infection of mice with *Candida albicans*. *J Infect Dis.* 2002a; 185:1833–1837. [PubMed: 12085336]
- Aratani Y, Kura F, Watanabe H, Akagawa H, Takano Y, Suzuki K, et al. Differential host susceptibility to pulmonary infections with bacteria and fungi in mice deficient in myeloperoxidase. *J Infect Dis.* 2000; 182:1276–1279. [PubMed: 10979934]
- Aratani Y, Kura F, Watanabe H, Akagawa Y, Takano Y, Suzuki K, et al. Relative contributions of myeloperoxidase and NADPH-oxidase to the early host defense against pulmonary infections with *Candida albicans* and *Aspergillus fumigatus*. *Med Mycol.* 2002b; 40:557–563. [PubMed: 12521119]
- Arnhold J, Monzani E, Furtmüller PG, Zederbauer M, Casella L, Obinger C. Kinetics and thermodynamics of halide and nitrite oxidation by mammalian heme peroxidases. *Eur J Inorg Chem.* 2006; 19:3801–3811.
- Brovkovich V, Gao XP, Ong E, Brovkovich S, Brennan ML, Su X, et al. Augmented inducible nitric oxide synthase expression and increased NO production reduce sepsis-induced lung injury and mortality in myeloperoxidase-null mice. *Am J Physiol Lung Cell Mol Physiol.* 2008; 295:L96–L103. [PubMed: 18424617]
- Chapman ALP, Hampton MB, Senthilmohan R, Winterbourn CC, Kettle AJ. Chlorination of bacterial and neutrophil proteins during phagocytosis and killing of *Staphylococcus aureus*. *J Biol Chem.* 2002; 277:9757–9762. [PubMed: 11733505]
- Clark RA. Extracellular effects of the myeloperoxidase-hydrogen peroxide-halide system. *Advances in Inflammation Research.* 1983; 5:107–146.
- Clark RA, Borregaard N. Neutrophils autoinactivate secretory products by myeloperoxidase-catalyzed oxidation. *Blood.* 1985; 65:375–381. [PubMed: 2981586]
- Clark RA, Klebanoff SJ. Chemotactic factor inactivation by the myeloperoxidase-hydrogen peroxide-halide system. An inflammatory control mechanism. *J Clin Invest.* 1979; 64:913–920. [PubMed: 225353]
- Clark RA, Stone PJ, El Hag A, Calore JD, Franzblau C. Myeloperoxidase-catalyzed inactivation of α 1-protease inhibitor by human neutrophils. *J Biol Chem.* 1981; 256:3348–3353. [PubMed: 6162845]
- Clark RA, Szot S, Venkatasubramanian K, Schiffmann E. Chemotactic factor inactivation by myeloperoxidase-mediated oxidation of methionine. *J Immunol.* 1980; 124:2020–2026. [PubMed: 6245132]
- Decleva E, Menegazzi R, Busetto S, Patriarca P, Dri P. Common methodology is inadequate for studies on the microbicidal activity of neutrophils. *J Leuk Biol.* 2006; 79:87–94.
- DeCoursey TE. Voltage-gated proton channels: molecular biology, physiology, and pathophysiology of the H(V) family. *Physiol Rev.* 2013; 93:599–652. [PubMed: 23589829]
- Dunford, HB. *Heme Peroxidases*. New York: Wiley-VCH; 1999.
- Dyrbukt JM, Bishop C, Brooks WM, Thong B, Eriksson H, Kettle AJ. A sensitive and selective assay for chloramine production by myeloperoxidase. *Free Radic Biol Med.* 2005; 39:1468–1477. [PubMed: 16274882]
- Eisenhauer PB, Lehrer RI. Mouse neutrophils lack defensins. *Infect Immun.* 1992; 60:3446–3447. [PubMed: 1639513]

- Elzanowska H, Wolcott RG, Hannum DM, Hurst JK. Bactericidal properties of hydrogen peroxide and copper or iron-containing complexes in relation to leukocyte function. *Free Radic Biol Med*. 1995; 18:437–449. [PubMed: 9101234]
- Espey MG, Xavier S, Thomas DD, Miranda KM, Wink DA. Direct real-time evaluation of nitration with green fluorescent protein in solution and within human cells reveals the impact of nitrogen dioxide vs. peroxynitrite mechanisms. *Proc Nat Acad Sci USA*. 2002; 99:3481–3486. [PubMed: 11904413]
- Foerder CA, Shapiro BM. Release of ovoperoxidase from sea urchin eggs hardens the fertilization membrane with tyrosine crosslinks. *Proc Natl Acad Sci U S A*. 1977; 74:4214–4218. [PubMed: 270665]
- Footo CS, Goynes TE, Lehrer RI. Assessment of chlorination by human neutrophils. *Nature*. 1983; 301:715–726. [PubMed: 6828155]
- Fu X, Kassim SY, Parks WC, Heinecke JW. Hypochlorous acid oxygenates the cysteine switch domain of pro-matrilysin (MMP-7). *J Biol Chem*. 2001; 276:41279–41287. [PubMed: 11533038]
- Fu X, Kassim SY, Parks WC, Heinecke JW. Hypochlorous acid generated by myeloperoxidase modifies adjacent tryptophan and glycine residues in the catalytic domain of matrix metalloproteinase-7 (matrilysin). *J Biol Chem*. 2003; 278:28403–28409. [PubMed: 12759346]
- Furtmüller PG, Arnhold J, Jantschko W, Pichler H, Obinger C. Redox properties of the couples compound I/compound II and compound II/native enzyme of human myeloperoxidase. *Biochem Biophys Res Commun*. 2003; 301:551–557. [PubMed: 12565898]
- Furtmüller PG, Burner U, Obinger C. Reaction of myeloperoxidase compound I with chloride, bromide, iodide, and thiocyanate. *Biochemistry*. 1998; 37:17923–17930. [PubMed: 9922160]
- Gaut JP, Yeh GC, Tran HD, Byun J, Henderson JP, Richter GM, et al. Neutrophils employ the myeloperoxidase system to generate antimicrobial brominating and chlorinating oxidants during sepsis. *Proc Nat Acad Sci USA*. 2001; 98:11961–11966. [PubMed: 11593004]
- Grignaschi VJ, Sperperato AM, Etcheverry MJ, Macario AJ. An nuevo cuadro citoquímico: Negatividad espontánea de las reacciones de peroxidación, oxidación y lípidos en la progenia neutrofilia y en los monocitos de dos hermanos. *Rev Assoc Med Argent*. 1963; 77:218–221.
- Haegens A, Heeringa P, van Suylen RJ, Steele C, Aratani Y, O'Donoghue RJJ, et al. Myeloperoxidase deficiency attenuates lipopolysaccharide-induced acute lung inflammation and subsequent cytokine and chemokine production. *J Immunol*. 2009; 182:7990–7996. [PubMed: 19494324]
- Hampton MB, Kettle AJ, Winterbourn CC. Involvement of superoxide and myeloperoxidase in oxygen-dependent killing of *Staphylococcus aureus* by neutrophils. *Infect Immun*. 1996; 64:3512–3517. [PubMed: 8751892]
- Harrison JE, Schultz J. Studies on the chlorinating activity of myeloperoxidase. *J Biol Chem*. 1976; 251:1371–1374. [PubMed: 176150]
- Hawkins CL, Davies MJ. Inactivation of protease inhibitors and lysozyme by hypochlorous acid: role of side-chain oxidation and protein unfolding in loss of biological function. *Chem Res Toxicol*. 2005; 18:1600–1610. [PubMed: 16533025]
- Hawkins CL, Pattison DI, Davies MJ. Hypochlorite-induced oxidation of amino acids, peptides, and proteins. *Amino Acids*. 2003; 25:259–274. [PubMed: 14661089]
- Hirche TO, Gaut JP, Heinecke JW, Belaaouaj A. Myeloperoxidase plays critical roles in killing *Klebsiella pneumoniae* and inactivating neutrophil elastase: effects on host defense. *J Immunol*. 2005; 174:1557–1565. [PubMed: 15661916]
- Holland SM. Chronic granulomatous disease. *Hematol Oncol Clin North Am*. 2013; 27:89–99. [PubMed: 23351990]
- Hurst JK. What really happens in the neutrophil phagosome? *Free Radic Biol Med*. 2012; 53:508–520. [PubMed: 22609248]
- Hurst JK, Barrette WC Jr. Leukocyte oxygen activation and microbicidal oxidative toxins. *CRC Crit Rev Biochem Mol Biol*. 1989; 24:271–328. [PubMed: 2548810]
- Imlay JA. Pathways of oxidative damage. *Annu Rev Microbiol*. 2003; 57:395–418. [PubMed: 14527285]
- Jandl RC, Andre S, Borges-DuBois JL, Kipnes RS, McMurrich BJ, Babior BM. Termination of the respiratory burst in human neutrophils. *J Clin Invest*. 1978; 61:1176–1185. [PubMed: 207730]

- Kitahara M, Eyre HJ, Simonian Y, Atkin CL, Hasstedt SJ. Hereditary myeloperoxidase deficiency. *Blood*. 1981; 57:888–893. [PubMed: 6260268]
- Klebanoff SJ. Iodination of bacteria; a bactericidal mechanism. *J Exp Med*. 1967; 126:1063–1078. [PubMed: 4964565]
- Klebanoff SJ. Myeloperoxidase-halide-hydrogen peroxide antibacterial system. *J Bacteriol*. 1968; 95:2131–2138. [PubMed: 4970226]
- Klebanoff, SJ. Myeloperoxidase-mediated antimicrobial systems and their role in leukocyte function. In: Schultz, J., editor. *Biochemistry of the Phagocytic Process*. North-Holland Publishing Company; 1970a. p. 89–110.
- Klebanoff SJ. Myeloperoxidase: contribution to the microbicidal activity of intact leukocytes. *Science*. 1970b; 169:1095–1097. [PubMed: 4988715]
- Klebanoff SJ. Myeloperoxidase: friend and foe. *J Leuk Biol*. 2005; 77:598–625.
- Klebanoff SJ, Hamon CB. Role of myeloperoxidase-mediated antimicrobial systems in intact leukocytes. *J Retic Soc*. 1972; 12:170–196.
- Klebanoff SJ, Kettle AJ, Rosen H, Winterbourn CC, Nauseef WM. Myeloperoxidase: a front-line defender against phagocytosed microorganisms. *J Leuk Biol*. 2013; 93:185–198.
- Klebanoff SJ, Pincus SH. Hydrogen peroxide utilization in myeloperoxidase-deficient leukocytes: a possible microbicidal control mechanism. *J Clin Invest*. 1971; 50:2226–2229. [PubMed: 5116211]
- Klebanoff, SJC.; RA. *The Neutrophil: Function and Clinical Disorders*. Amsterdam: Elsevier/North Holland Biomedical Press; 1978. p. 810
- Kobayashi M, Tanaka T, Usui T. Inactivation of lysosomal enzymes by the respiratory burst of polymorphonuclear leukocytes. *J Lab Clin Med*. 1982; 100:896–907. [PubMed: 6292313]
- Koch C. Neutrophil granulocyte function *in vitro*. *Acta Pathol Microbiol Scand B Microbiol Immunol*. 1974; 82:127–135. [PubMed: 4275242]
- Kutter D. Prevalence of myeloperoxidase deficiency: population studies using Bayer-Technicon automated hematology. *J Mol Med*. 1998; 76:669–675. [PubMed: 9766844]
- Kutter D, Al-Haidari K, Thoma J. Myeloperoxidase deficiency: simple methods for its diagnosis and significance of different forms. *Klinisches Labor*. 1994; 40:342–346.
- Lehrer RI, Cline MJ. Leukocyte myeloperoxidase deficiency and disseminated candidiasis: the role of myeloperoxidase in resistance to *Candida* infection. *J Clin Invest*. 1969a; 48:1478–1488. [PubMed: 5796360]
- Lehrer RI, Hanifin J, Cline MJ. Defective bactericidal activity in myeloperoxidase-deficient human neutrophils. *Nature*. 1969b; 223:78–79. [PubMed: 4978299]
- Lymar SV, Hurst JK. Role of compartmentation in promoting toxicity of leukocyte-generated strong oxidants. *Chem Res Toxicol*. 1995; 8:833–840. [PubMed: 7492732]
- Malle E, Buch T, Grone HJ. Myeloperoxidase in kidney disease. *Kidney Int*. 2003; 64:1956–1967. [PubMed: 14633118]
- Marquez LA, Huang JT, Dunford HB. Spectral and kinetic studies on the formation of myeloperoxidase compounds I and II: Roles of hydrogen peroxide and superoxide. *Biochemistry*. 1994; 33:1447–1454. [PubMed: 8312264]
- Mestas J, Hughes CCW. Of mice and not men: differences between mouse and human immunology. *J Immunol*. 2004; 172:2731–2738. [PubMed: 14978070]
- Moreland JG, Davis AP, Bailey G, Nauseef WM, Lamb FS. Anion channels, including CLC-3, are required for normal neutrophil oxidative function, phagocytosis, and transendothelial migration. *J Biol Chem*. 2006; 281:12277–12288. [PubMed: 16522634]
- Nauseef WM. How human neutrophils kill and degrade microbes: an integrated view. *Immunol Rev*. 2007; 219:88–102. [PubMed: 17850484]
- Nauseef WM, Metcalf JA, Root RK. Role of myeloperoxidase in the respiratory burst of human neutrophils. *Blood*. 1983; 61:483–491. [PubMed: 6297637]
- Nicholls SJ, Hazen SL. Myeloperoxidase, modified lipoproteins, and atherogenesis. *J Lipid Res*. 2009; 50:S346–S351. [PubMed: 19091698]
- Nunoi H, Kohi F, Kajiwara H, Suzuki K. Prevalence of inherited myeloperoxidase deficiency in Japan. *Microbiol Immunol*. 2003; 47:527–531. [PubMed: 12953846]

- Nussbaum C, Klinke A, Adam M, Baldus S, Sperandio M. Myeloperoxidase - a leukocyte-derived protagonist of inflammation and cardiovascular disease. *Antioxidants & Redox Signaling*. 2013; 18:692–713. [PubMed: 22823200]
- Odeberg H, Olsson I. Microbicidal mechanisms of human granulocytes: synergistic effects of granulocyte elastase and myeloperoxidase or chymotrypsin-like cationic protein. *Infect Immun*. 1976; 14:1276–1283. [PubMed: 12111]
- Painter RG, Bonvillain RW, Valentine VG, Laombard GA, Laplace SG, Nauseef WM, Wang G. The role of chloride anion and CFTR in killing of *Pseudomonas aeruginosa* by normal and CF neutrophils. *J Leuk Biol*. 2008; 83:1345–1353.
- Painter RG, Marrero L, Lombard GA, Valentine VG, Nauseef WM, Wang G. CFTR-mediated halide transport in phagosomes of human neutrophils. *J Leuk Biol*. 2010; 87:933–942.
- Painter RG, Valentine VG, Lanson NA Jr, Leidal K, Zhang O, Lombard G, et al. CFTR expression in human neutrophils and the phagolysosomal chlorination defect in cystic fibrosis. *Biochemistry*. 2006; 45:10260–10269. [PubMed: 16922501]
- Palazzolo AM, Suquet C, Konkel ME, Hurst JK. Green fluorescent protein-expressing *Escherichia coli* as a selective probe for HOCl generation within neutrophils. *Biochemistry*. 2005; 44:6910–6919. [PubMed: 15865436]
- Pang YY, Schwartz J, Bloomberg S, Boyd JM, Horswill AR, Nauseef WM. Methionine Sulfoxide Reductases Protect against Oxidative Stress in *Staphylococcus aureus* Encountering Exogenous Oxidants and Human Neutrophils. *J Innate Immun*. 2014; 6:353–364. [PubMed: 24247266]
- Park S, You X, Imlay JA. Substantial DNA damage from submicromolar intracellular hydrogen peroxide detected in Hpx- mutants of *Escherichia coli*. *Proc Nat Acad Sci USA*. 2005; 102:9317–9322. [PubMed: 15967999]
- Pattison DI, Davies MJ. Absolute rate constants for the reactions of hypochlorous acid with protein side chains and peptide bonds. *Chem Res Toxicol*. 2001; 14:1453–1464. [PubMed: 11599938]
- Pattison DI, Davies MJ. Reactions of myeloperoxidase-derived oxidants with biological substrates: gaining chemical insight into human inflammatory diseases. *Curr Med Chem*. 2006; 13:3271–3290. [PubMed: 17168851]
- Rausch PG, Moore TG. Granule enzymes of polymorphonuclear neutrophils: a phylogenetic comparison. *Blood*. 1975; 46:913–919. [PubMed: 173439]
- Rittirsch D, Hoesel LM, Ward PA. The disconnect between animal models of sepsis and human sepsis. *J Leuk Biol*. 2007; 81:137–143.
- Rorvig S, Ostergaard O, Heegaard NH, Borregaard N. Proteome profiling of human neutrophil granule subsets, secretory vesicles, and cell membrane: correlation with transcriptome profiling of neutrophil precursors. *J Leukoc Biol*. 2013; 94:711–721. [PubMed: 23650620]
- Rosen H, Klebanoff SJ. Chemiluminescence and superoxide production by myeloperoxidase-deficient leukocytes. *J Clin Invest*. 1976; 58:50–60. [PubMed: 180060]
- Rosen H, Klebanoff SJ, Wang Y, Brot N, Heinecke JW, Fu X. Methionine oxidation contributes to bacterial killing by the myeloperoxidase system of neutrophils. *Proc Nat Acad Sci USA*. 2009; 106:18686–18691. [PubMed: 19833874]
- Rosen H, Michel BR. Redundant contribution of myeloperoxidase-dependent systems to neutrophil-mediated killing of *Escherichia coli*. *Infect Immun*. 1997; 65:4173–4178. [PubMed: 9317024]
- Ruf J, Carayon P. Structural and functional aspects of thyroid peroxidase. *Arch Biochem Biophys*. 2006; 445:269–277. [PubMed: 16098474]
- Schultz J, Kaminker K. Myeloperoxidase of the leucocyte of normal human blood. *Arch Biochem Biophys*. 1962; 96:465–467. [PubMed: 13909511]
- Schwartz J, Leidal KG, Femling JK, Weiss JP, Nauseef WM. Neutrophil bleaching of GFP-expressing staphylococci: probing the intraphagosomal fate of individual bacteria. *J Immunology*. 2009; 183:2632–2641. [PubMed: 19620311]
- Shao B, Belaouaj A, Verlinde CLMJ, Fu X, Heinecke JW. Methionine sulfoxide and proteolytic cleavage contribute to the inactivation of cathepsin G by hypochlorous acid. An oxidative mechanism for regulation of serine proteinases by myeloperoxidase. *J Biol Chem*. 2005; 280:29311–29321. [PubMed: 15967795]

- Shao B, Pennathur S, Heinecke JW. Myeloperoxidase targets apolipoprotein A01, the major high density lipoprotein protein, for site-specific oxidation in human atherosclerotic lesions. *J Biol Chem.* 2012; 287:6375–6386. [PubMed: 22219194]
- Staudinger BJ, Oberdoerster MA, Lewis PJ, Rosen H. mRNA expression profiles for *Escherichia coli* ingested by normal and phagocyte oxidase-deficient human neutrophils. *J Clin Invest.* 2002; 110:1151–1163. [PubMed: 12393851]
- Summers FA, Morgan PE, Davies MJ, Hawkins CL. Identification of plasma proteins that are susceptible to thiol oxidation by hypochlorous acid and N-chloramines. *Chem Res Toxicol.* 2008; 21:1832–1840. [PubMed: 18698849]
- Sun YT, Shieh CC, Delpire E, Shen MR. K^+ - Cl^- cotransport mediates the bactericidal activity of neutrophils by regulating NADPH oxidase activation. *J Physiol.* 2012; 590:3231–3243. [PubMed: 22526882]
- Takeuchi K, Umeki Y, Matsumoto N, Yamamoto K, Yoshida M, Suzuki K, Aratani Y. Severe neutrophil-mediated lung inflammation in myeloperoxidase-deficient mice exposed to zymosan. *Inflamm Res.* 2012; 61:197–205. [PubMed: 22116298]
- Theorell H, Åkeson. Highly purified milk peroxidase. *Arkiv för Kemi, Mineralogi, och Geologi.* 1943; 17:1–6.
- Thomas EL. Myeloperoxidase-hydrogen peroxide-chloride antimicrobial system: effect of exogenous amines on antibacterial action against *Escherichia coli*. *Infect Immun.* 1979a; 25:110–116. [PubMed: 39030]
- Thomas EL. Myeloperoxidase, hydrogen peroxide, chloride antimicrobial system: nitrogen-chlorine derivatives of bacterial components in bactericidal action against *Escherichia coli*. *Infect Immun.* 1979b; 23:522–531. [PubMed: 217834]
- Thomas EL, Grisham MB, Jefferson MM. Myeloperoxidase-dependent effect of amines on functions of isolated neutrophils. *J Clin Invest.* 1983; 72:441–454. [PubMed: 6308055]
- Thomas EL, Grisham MB, Jefferson MM. Cytotoxicity of chloramines. *Methods Enzymol.* 1986; 132:585–593. [PubMed: 3547027]
- Thomas EL, Jefferson MM, Grisham MB. Myeloperoxidase-catalyzed incorporation of amines into proteins: role of hypochlorous acid and dichloramines. *Biochemistry.* 1982; 21:6299–6308. [PubMed: 6295461]
- Undritz E. Die Alius-Grignaschi-Anomalie: der erblich konstitutionelle Peroxydasedefekt der Neutrophilien und Monozyten. *Blut.* 1966; 14:129–136. [PubMed: 5957405]
- Undurti A, Huang Y, Lupica JA, Smith JD, DiDonato JA, Hazen SL. Modification of high density lipoprotein by myeloperoxidase generates a pro-inflammatory particle. *J Biol Chem.* 2009; 284:30825–30835. [PubMed: 19726691]
- Vissers MCM, Winterbourn CC. Oxidative damage to fibronectin. I The effects of the neutrophil myeloperoxidase system and HOCl. *Arch Biochem Biophys.* 1991; 285:53–59. [PubMed: 1846732]
- Voetman AA, Weening RS, Hamers MN, Meerhof LJ, Bot AA, Roos D. Phagocytosing human neutrophils inactivate their own granular enzymes. *J Clin Invest.* 1981; 67:1541–1549. [PubMed: 7229038]
- Weiss SJ, Klein R, Slivka A, Wei M. Chlorination of taurine by human neutrophils. Evidence for hypochlorous acid generation. *J Clin Invest.* 1982; 70:598–607. [PubMed: 6286728]
- Weiss SJ, Peppin G, Ortiz X, Ragsdale C, Test ST. Oxidative autoactivation of latent collagenase by human neutrophils. *Science.* 1985; 227:747–749. [PubMed: 2982211]
- Winter J, Ilbert M, Graf PCF, Özcelik D, Jakob U. Bleach activates a redox-regulated chaperone by oxidative protein unfolding. *Cell.* 2008; 135:691–701. [PubMed: 19013278]
- Winterbourn CC, Hampton MB, Livesey JH, Kettle AJ. Modeling the reactions of superoxide and myeloperoxidase in the neutrophil phagosome. *J Biol Chem.* 2006; 281:39860–39869. [PubMed: 17074761]
- Winterbourn CC, Kettle AJ. Redox reactions and microbial killing in the neutrophil phagosome. *Antioxid Redox Signal.* 2013; 18:642–660. [PubMed: 22881869]

- Zamocky M, Jakopitsch C, Furtmüller PG, Dunand C, Obinger C. The peroxidase-cyclooxygenase superfamily: reconstructed evolution of critical enzymes of the innate immune system. *Proteins:structure, function, and genetics*. 2008; 72:589–605.
- Zederbauer M, Furtmüller PG, Brogioni S, Jakopitsch C, Smulevich G, Obinger C. Heme to protein linkages in mammalian peroxidases: impact on spectroscopic, redox, and catalytic properties. *Nature Product Reports*. 2007; 24:571–584.
- Zgliczynski JM. Characteristics of MPO from neutrophils and other peroxidases from different cells. In: Sbarra, AJ.; Strauss, RR., editors. *The Reticuloendothelial System. A Comprehensive Treatise. Volume 2: Biochemistry and Metabolism*. New York: Plenum Press; 1980. p. 255-278.
- Zgliczynski JM, Stelmaszynska T, Domanski J, Ostrowski W. Chloramines as intermediates of oxidative reaction of amino acids by myeloperoxidase. *Biochem Biophys Acta*. 1971; 235:419–424. [PubMed: 4378090]
- Zgliczynski JM, Stelmaszynska T, Ostrowski W, Naskalski J, Sznajd J. Myeloperoxidase of human leukaemic leucocytes. Oxidation of amino acids in the presence of hydrogen peroxide. *Eur J Biochem*. 1968; 4:540–547. [PubMed: 5242086]
- Zgliczynski JM, Stelmaszynska T. Hydrogen cyanide and cyanogen chloride formation by the myeloperoxidase-H₂O₂-Cl⁻ system. *Biochim Biophys Acta*. 1979; 567:309–314. [PubMed: 36154]
- Zhou Y, Song K, Painter RG, Aiken M, Reiser J, Stanton BA, et al. Cystic fibrosis transmembrane conductance regulator recruitment to phagosomes in neutrophils. *J Innate Immun*. 2013; 5:219–230. [PubMed: 23486169]




Figure. MPO-dependent events in human neutrophil phagosomes

Concomitant with phagocytosis, human neutrophils assemble and activate the NADPH oxidase and recruit granules to fuse with nascent phagosomes. The NADPH oxidase transfers electrons into the phagosome, generating superoxide anion ($O_2^{\cdot -}$) and hydrogen peroxide (H_2O_2) from molecular oxygen. The voltage-gated proton channel Hv1 provides the bulk of charge compensation necessitated by electron transfer from cytoplasm, thereby allowing sustained generation of oxidants in close proximity to the ingested microbe. Chloride (Cl^-) from the neutrophil cytoplasm enters phagosomes predominantly via cystic fibrosis conductance regulator (CFTR), although ClC3 and KCC3 support some Cl^- influx. MPO, provided by azurophilic granules, and H_2O_2 react to yield Compound I (Cpd I), which catalyzes the two electron oxidation of Cl^- to Cl^+ to produce hypochlorous acid (HOCl) or bleach. HOCl reacts with peptides and proteins from host and microbe to generate an array of products, including monochloramines (NH_2Cl), protein chloramines (PNHCl), which can then decompose to form aldehydes. Granule proteins, both native and oxidant-modified, HOCl and its derivatives attack microbial targets, with the outcome dependent both on the number and accessibility of vulnerable targets as well on the defensive responses specific to the ingested organism.